


FKSG80 / GPR81 Rabbit anti-Human Polyclonal (Cytoplasmic Domain) (Unconjugated) Antibody - LS-A2093 - LSBio	
CatalogID:	LS-A2093
Target:	G protein-coupled receptor 81 (HCAR1)
Synonyms:	HCAR1 Antibody, FKSG80 Antibody, G protein-coupled receptor 81 Antibody, G-protein coupled receptor 104 Antibody, G protein-coupled receptor 104 Antibody, G-protein coupled receptor 81 Antibody, GPR81 Antibody, GPR104 Antibody, LACR1 Antibody, HCA1 Antibody, TA-GPCR Antibody, Lactate receptor 1 Antibody
Family / Subfamily:	GPCR / Chemokine
Host	HCAR1 antibody was produced in Rabbit
Clonality:	Polyclonal
Modifications:	Unconjugated (Unconjugated)
Immunogen Species:	FKSG80 / GPR81 antibody was raised against Human
Immunogen:	FKSG80 / GPR81 antibody was raised against synthetic 17 amino acid peptide from 3rd cytoplasmic domain of human GPR81. Percent identity with other species by BLAST analysis: Human, Gorilla, Monkey (100%); Bat (94%); Marmoset, Platypus (88%); Mouse, Hamster (82%).
Specificity:	Human GPR81. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Epitope:	Cytoplasmic Domain
Reactivity:	Human
Predicted Reactivity:	Bat, Gorilla, Monkey
Purification:	Immunoaffinity Chromatography
Presentation:	PBS, 0.1% sodium azide.
Recommended Storage:	Long term: -70°C; Short term: +4°C
Usage Summary:	Immunohistochemistry: LS-A2093 was validated for use in immunohistochemistry on a panel of 21 formalin-fixed, paraffin-embedded (FFPE) human tissues after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with the primary antibody, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen. The stained slides were evaluated by a pathologist to confirm staining specificity. The optimal working concentration for LS-A2093 was determined to be 11 ug/ml.
Uses:	IHC-P (11 μg/ml) (Optimal dilution to be determined by the researcher)
Size:	50 μg
Concentration:	1 mg/ml



Anti-GPR81 antibody LS-A2093 IHC staining of human colon, epithelium. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

Requested From: United States

Laboratory Reagent For In Vitro Research Use Only
Not for resale without prior written consent from LifeSpan BioSciences, Inc.
Created on 5/24/2013
© 2013 LifeSpan BioSciences